

### **Product Datasheet**



# **DYKDDDDK-Tag Mouse Monoclonal Antibody**

**Catalog #: EAB10043** 

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IP, IF/ICC	N/A	Species independent

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:10000-50000
IP(Immunoprecipitation) 1:100-500
IF/ICC(Immunofluorescence/Immunocytochemistry) 1:500-3000

### **Product Information**

**Conjugate** Unconjugate

Specificity

DYKDDDDK-Tag Mouse Monoclonal Antibody detects endogenous levels of DYKDDDDK-Tag

fusion protein.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

**Formulation** In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks.

UniProt ID N/A
Entrez-Gene ID N/A

## **Product Description**

DYKDDDDK-Tag (Flag-tag) is a small component of an epitope which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein. It has been used extensively as a general epitope tag in expression vectors. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits. A Flag-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against the studied protein, adding a Flag-tag to this protein allows one to follow the protein with an antibody against the Flag sequence.