

## PI3 Kinase p85 $\alpha$ Rabbit Polyclonal Antibody

### Catalog #: EAB10100

| Host/Isotype | Clonality  | Applications             | MW (kDa) | Reactivity             |
|--------------|------------|--------------------------|----------|------------------------|
| Rabbit IgG   | Polyclonal | WB, IHC-P, IF/ICC, ELISA | 84       | Human, Mouse, Rat, Pig |

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

|                                                        |              |
|--------------------------------------------------------|--------------|
| <b>WB</b> (Western Blotting)                           | 1:500-2000   |
| <b>IHC-P</b> (Immunohistochemistry-Paraffin)           | 1:50-300     |
| <b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry) | 1:50-300     |
| <b>ELISA</b> (Enzyme-linked Immunosorbent Assay)       | 1:5000-20000 |

### Product Information

|                       |                                                                                                                                           |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Conjugate</b>      | Unconjugate                                                                                                                               |
| <b>Specificity</b>    | PI3 Kinase p85 $\alpha$ Rabbit Polyclonal Antibody detects endogenous levels of PI3 Kinase p85 $\alpha$ protein.                          |
| <b>Purification</b>   | Affinity purification                                                                                                                     |
| <b>Concentration</b>  | 1mg/ml                                                                                                                                    |
| <b>Format</b>         | Liquid                                                                                                                                    |
| <b>Formulation</b>    | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol                                                                  |
| <b>Shipping</b>       | Gel Pack                                                                                                                                  |
| <b>Storage</b>        | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles.<br>Aliquots may be stored at +4°C for 1-2 weeks |
| <b>UniProt ID</b>     | <a href="#">P27986</a>                                                                                                                    |
| <b>Entrez-Gene Id</b> | <a href="#">5295</a>                                                                                                                      |

### Product Description

Phosphatidylinositol 3-kinase (PI 3-kinase) is composed of p85 and p110 subunits. p85 lacks PI 3-kinase activity and acts as an adapter, coupling p110 to activated protein tyrosine kinase. Two forms of p85 have been described (p85 $\alpha$  and p85 $\beta$ ), each possessing one SH3 and two SH2 domains. Various p110 isoforms have been identified. p110 $\alpha$  and p110 $\beta$  interact with p85 $\alpha$ , and p110 $\alpha$  has also been shown to interact with p85 $\beta$  in vitro. p110 $\delta$  expression is restricted to white blood cells. It has been shown to bind p85  $\alpha$  and  $\beta$ , but it apparently does not phosphorylate these subunits. p110 $\delta$  seems to have the capacity to autophosphorylate. p110 $\gamma$  does not interact with the p85 subunits. It has been shown to be activated by  $\alpha$  and  $\beta\gamma$  heterotrimeric G proteins.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**