

Order:

Web:

Phospho-HER2/ErbB2 (Tyr877) Rabbit Polyclonal Antibody

Catalog #: EAB10252

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IP, IHC-P, IF, ELISA	137	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

IP(Immunoprecipitation)1:20-200IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000	WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300		
IF(Immunofluorescence) 1:50-300		
	(, , , , , , , , , , , , , , , , , , ,	
ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000	IF(Immunofluorescence)	1:50-300
	ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Phospho-HER2/ErbB2 (Tyr877) Rabbit Polyclonal Antibody detects endogenous levels of HER2/ErbB2 only when phosphorylated at Tyr877.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>P04626</u>
Entrez-Gene Id	2064

Product Description

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA