

#### **Product Datasheet**

Order: order@ebiocell.com

TEL: (540)808-3925 tech@ebiocell.com

Supprt: Web: www.ebiocell.com

# Phospho-FoxO3A (Ser413) Rabbit Polyclonal Antibody

Catalog #: EAB10344

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB	71	Human, Mouse, Rat

### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user. 1:500-2000

**WB**(Western Blotting)

## **Product Information**

Conjugate Unconjugate

Phospho-FoxO3A (Ser413) Rabbit Polyclonal Antibody detects endogenous levels of FoxO3A Specificity

protein only when phosphorylated at Ser413.

**Purification** Affinity purification

Concentration 1mg/ml **Format** Liquid

**Formulation** In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping

Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Storage

Aliquots may be stored at +4°C for 1-2 weeks

**UniProt ID** O43524 **Entrez-Gene Id** 2309

## **Product Description**

FOXO3 (forkhead box O3) or FOXO3A, also known as FKHRL1 (forkhead in rhabdomyosarcoma-like 1), is a 673 amino acid transcriptional activator that belongs to the FKHR subfamily of forkhead transcription factors. Transcriptional activation of FKHR proteins is regulated by the serine/threonine kinase Akt1, which phosphorylates FKHRL1 at Threonine 32 and Serine 253. Phosphorylation by Akt1 negatively regulates FKHRL1 by promoting its export from the nucleus. Phosphorylated FKHRL1 associates with 14-3-3 proteins and this complex is retained in the cytoplasm. Growth factor withdrawal stimulates FKHRL1 dephosphorylation and nuclear translocation, leading to FKHRinduced gene-specific transcriptional activation. Within the nucleus, dephosphorylated FKHRL1 triggers apoptosis by inducing the expression of genes that are critical for cell death.