

Phospho- α -Synuclein (Ser129) Rabbit Polyclonal Antibody

Catalog #: EAB10555

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|----------------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC-P, IF, ELISA | 14 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|--------------|
| WB (Western Blotting) | 1:500-2000 |
| IHC-P (Immunohistochemistry-Paraffin) | 1:50-300 |
| IF (Immunofluorescence) | 1:50-300 |
| ELISA (Enzyme-linked Immunosorbent Assay) | 1:5000-20000 |

Product Information

| | |
|-----------------------|---|
| Conjugate | Unconjugate |
| Specificity | Phospho- α -Synuclein (Ser129) Rabbit Polyclonal Antibody detects endogenous levels of α -Synuclein protein only when phosphorylated at Ser129. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | P37840 |
| Entrez-Gene Id | 6622 |

Product Description

Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Alternatively spliced transcripts encoding different isoforms have been identified for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures