

Phospho-TAK1 (Thr187) Rabbit Polyclonal Antibody

Catalog #: EAB10593

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	67	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF(Immunofluorescence)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Unconjugate	
Phospho-TAK1 (Thr187) Rabbit Polyclonal Antibody detects endogenous levels of TAK1 only when phosphorylated at Thr187.	
Affinity purification	
1mg/ml	
Liquid	
In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol	
Gel Pack	
Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks	
<u>O43318</u>	
<u>6885</u>	

Product Description

The protein encoded by this gene is a member of the serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA