

Order:

Supprt:

Web:

Phospho-CDK1/Cdc2 (Tyr15) Rabbit Polyclonal Antibody

Catalog #: EAB10606

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, ELISA	34	Human, Mouse, Rat, Monkey

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Phospho-CDK1/Cdc2 (Tyr15) Rabbit Polyclonal Antibody detects endogenous levels of CDK1/Cdc2 protein only when phosphorylated at Tyr15.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>P06493</u>
Entrez-Gene Id	<u>983</u>

Product Description

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associate with this protein and function as regulatory subunits. The kinase activity of this protein also controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA