

## Phospho-Histone H2A.X (Thr120) Rabbit Polyclonal Antibody

### Catalog #: EAB10791

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB	15	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**WB**(Western Blotting) 1:500-2000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Phospho-Histone H2A.X (Thr120) Rabbit Polyclonal Antibody detects endogenous levels of histone H2A.X protein only when phosphorylated at Thr120.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P16104</a>
<b>Entrez-Gene Id</b>	<a href="#">3014</a>

### Product Description

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**