

## Phospho-HP1 $\gamma$ (Ser83) Rabbit Polyclonal Antibody

### Catalog #: EAB10799

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB	21	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**WB**(Western Blotting) 1:500-2000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Phospho-HP1 $\gamma$ (Ser83) Rabbit Polyclonal Antibody detects endogenous levels of HP1 $\gamma$ protein only when phosphorylated on Ser83.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">Q13185</a>
<b>Entrez-Gene Id</b>	<a href="#">11335</a>

### Product Description

At the nuclear envelope, the nuclear lamina and heterochromatin are adjacent to the inner nuclear membrane. The protein encoded by this gene binds DNA and is a component of heterochromatin. This protein also can bind lamin B receptor, an integral membrane protein found in the inner nuclear membrane. The dual binding functions of the encoded protein may explain the association of heterochromatin with the inner nuclear membrane. This protein binds histone H3 tails methylated at Lys-9 sites. This protein is also recruited to sites of ultraviolet-induced DNA damage and double-strand breaks. Two transcript variants encoding the same protein but differing in the 5' UTR, have been found for this gene.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**