

## MAPKAPK-3 Rabbit Polyclonal Antibody

### Catalog #: EAB11794

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	43	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-300
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-300
<b>ELISA</b> (Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	MAPKAPK-3 Rabbit Polyclonal Antibody detects endogenous levels of MAPKAPK-3 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">Q16644</a>
<b>Entrez-Gene ID</b>	<a href="#">7867</a>

### Product Description

MAPKAP kinase-3 also designated 3PK, MAPKAPK-3 is a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and cell differentiation. Alternate splicing results in multiple transcript variants that encode the same protein.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**