

PPAR δ Rabbit Polyclonal Antibody

Catalog #: EAB13059

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|--------------|
| Rabbit IgG | Polyclonal | WB, IF | 50 | Human, Mouse |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|------------------------|------------|
| WB(Western Blotting) | 1:500-2000 |
| IF(Immunofluorescence) | 1:50-300 |

Product Information

| | |
|----------------|---|
| Conjugate | Unconjugate |
| Specificity | PPAR δ Rabbit Polyclonal Antibody detects endogenous levels of PPAR δ protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | Q03181 |
| Entrez-Gene Id | 5467 |

Product Description

This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) family. The encoded protein is thought to function as an integrator of transcriptional repression and nuclear receptor signaling. It may inhibit the ligand-induced transcriptional activity of peroxisome proliferator activated receptors alpha and gamma, though evidence for this effect is inconsistent. Expression of this gene in colorectal cancer cells may be variable but is typically relatively low. Knockout studies in mice suggested a role for this protein in myelination of the corpus callosum, lipid metabolism, differentiation, and epidermal cell proliferation. Alternative splicing results in multiple transcript variants encoding distinct protein isoforms.

For Research Use Only. Not For Use In Diagnostic Procedures