

## OB-Cadherin Rabbit Polyclonal Antibody

### Catalog #: EAB13082

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, ELISA	88	Human, Mouse

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>ELISA</b> (Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	OB-Cadherin Rabbit Polyclonal Antibody detects endogenous levels of OB-Cadherin protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
<b>UniProt ID</b>	<a href="#">P55287</a>
<b>Entrez-Gene ID</b>	<a href="#">1009</a>

### Product Description

OB-cadherin (also designated cadherin 11, OSF-4, CDH11) is a type II classical cadherin from the cadherin superfamily, integral membrane proteins that mediate calcium-dependent cell-cell adhesion. Mature cadherin proteins are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. Type II (atypical) cadherins are defined based on their lack of a HAV cell adhesion recognition sequence specific to type I cadherins. Expression of this particular cadherin in osteoblastic cell lines, and its upregulation during differentiation, suggests a specific function in bone development and maintenance.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**