

Product Datasheet

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

Phospho-Topoisomerase IIα (Thr1343) Rabbit Polyclonal Antibody

Catalog #: EAB13199

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, ELISA	174	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000 ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000

Product Information

Conjugate Unconjugate

Specificity

Phospho-Topoisomerase IIα (Thr1343) Rabbit Polyclonal Antibody detects endogenous levels of

topoisomerase IIα protein only when phosphorylated at Ser1343.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P11388

 Entrez-Gene Id
 7153

Product Description

This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This nuclear enzyme is involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. It catalyzes the transient breaking and rejoining of two strands of duplex DNA which allows the strands to pass through one another, thus altering the topology of DNA. Two forms of this enzyme exist as likely products of a gene duplication event. The gene encoding this form, alpha, is localized to chromosome 17 and the beta gene is localized to chromosome 3. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also play a role in ataxia-telangiectasia.