

## XRCC2 Rabbit Polyclonal Antibody

### Catalog #: EAB14107

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	32	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-300
<b>IF</b> (Immunofluorescence)	1:50-300
<b>ELISA</b> (Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	XRCC2 Rabbit Polyclonal Antibody detects endogenous levels of XRCC2 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
<b>UniProt ID</b>	<a href="#">O43543</a>
<b>Entrez-Gene ID</b>	<a href="#">7516</a>

### Product Description

This gene encodes a member of the RecA/Rad51-related protein family that participates in homologous recombination to maintain chromosome stability and repair DNA damage. This gene is involved in the repair of DNA double-strand breaks by homologous recombination and it functionally complements Chinese hamster irs1, a repair-deficient mutant that exhibits hypersensitivity to a number of different DNA-damaging agents.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**