

Product Datasheet

Order: order@ebiocell.com

Supprt:

Web:

TEL: (540)808-3925 tech@ebiocell.com

www.ebiocell.com

Phospho-p70 S6 kinase (Thr421) Rabbit Polyclonal Antibody

Catalog #: EAB14781

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	59	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)1:500-2000IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000

Product Information

Conjugate Unconjugate

Specificity

Phospho-p70 S6 kinase (Thr421) Rabbit Polyclonal Antibody detects endogenous levels of p70

S6 kinase β protein only when phosphorylated at Thr421.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 Q9UBS0

 Entrez-Gene Id
 6199

Product Description

In studies to elucidate key regulatory pathways in signal transduction, several protein serine/threonine (Ser/Thr) kinases have been identified, including two distinct families of 40S ribosomal protein S6 Ser/Thr kinases present in somatic animal cells, designated p70 S6 kinase and p90 Rsk kinase. p90 Rsk kinase is maximally activated within minutes of addition of growth factors or phorbol ester to cultured cells followed by activation of p70 S6 kinase. Both enzymes are regulated by serine/threonine phosphorylation, suggesting that specific kinases may exist upstream in the signaling pathway that regulate these kinases. In fact, evidence suggests that one such family of activating enzymes includes the members of the ERK MAP kinase family. The ERK MAP kinases are, in turn, regulated by phosphorylation at threonine and tyrosine residues by a protein kinase designated MEK.