

KAP1/TIF1 β Rabbit Monoclonal Antibody

Catalog #: EAB21230

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P, IF/ICC, FC	89	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IHC-P (Immunohistochemistry-Paraffin)	1:50-200
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-200
FC (Flow Cytometry)	1:10-100

Product Information

Conjugate	Unconjugate
Specificity	KAP1/TIF1 β Rabbit Monoclonal Antibody detects endogenous levels of KAP1/TIF1 β protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	Q13263
Entrez-Gene ID	10155

Product Description

TIF1 β , for transcriptional intermediary factor 1-beta, also designated KAP1(for KRAB-associated protein 1), TF1 β and TRIM28 (for tripartite motif-containing 28), is a member of the tripartite motif family characterized by three zinc-binding domains, a RING finger, B-boxes and a coiled-coil domain. Like TIF1 α , TIF1 β contains both a Cys/His PHD (plant homeodomain) finger and bromodomain that form a cooperative unit required for transcriptional repression. TIF1 β mediates transcriptional control by interaction with the Kruppel-associated box (KRAB) repression domain found in many transcription factors and by binding DNA through its zinc finger. The human TIF1 β gene maps to human chromosome 19q13.4 and encodes an 835 amino acid nuclear protein.

For Reserch Use Only. Not For Use In Diagnostic Procedures