

### **Product Datasheet**

Order:

order@ebiocell.com TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

# McI-1 Rabbit Monoclonal Antibody

Catalog #: EAB21282

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC, FC	37	Human, Mouse, Rat

### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
IP(Immunoprecipitation)	1:10-100
IHC-P(Immunohistochemistry-Paraffin)	1:50-200
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-200
FC(Flow Cytometry)	1:10-100

#### **Product Information**

Conjugate Unconjugate

Specificity McI-1 Rabbit Monoclonal Antibody detects endogenous levels of McI-1 protein.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 Q07820

 Entrez-Gene Id
 4170

## **Product Description**

Mcl-1 (also designated MCL1 or Myeloid cell leukemia-1) is an anti-apoptotic protein, which is a member of the Bcl-2 family. B-cell CLL/lymphoma 2 (Bcl-2) blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal. Myeloid cell leukemia 1 (Mcl-1) shares sequence homology with Bcl-2 and further resembles Bcl-2 in that its expression promotes cell viability. p53 and Mcl-1 demonstrate opposing effects on mitochondrial apoptosis by mediating Bcl-2 antagonist killer (Bak) activity. Mcl-1 is an important and specific regulator that is necessary for the homeostasis of early hematopoietic progenitors. Glycogen synthase kinase 3 (GSK3) controls Mcl-1 stability, which has an effect on the regulation of apoptosis by growth factors, PI 3-kinase and AKT. Mice with a deficiency of the Mcl-1 protein show a significant reduction in B and T lymphocytes similar to the effects observed in IL-7- or IL-7R-deficient mice.