

JNK1/JNK2/JNK3 Rabbit Monoclonal Antibody

Catalog #: EAB21328

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IF/ICC	48, 53	Human, Mouse, Rat, Cow, Monkey, Zebrafish

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IP (Immunoprecipitation)	1:10-100
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-200

Product Information

Conjugate	Unconjugate
Specificity	JNK1/JNK2/JNK3 Rabbit Monoclonal Antibody detects endogenous levels of JNK1/JNK2/JNK3 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P45983 , P45984 , P53779
Entrez-Gene Id	5599 , 5601 , 5602

Product Description

JNK proteins phosphorylate and augment transcriptional activity of c-Jun. JNKs originate from three genes that yield 10 isoforms through alternative mRNA splicing, including JNK1a1, JNK1b1, JNK2a1, JNK2b1, and JNK3a1, which represent the p46 isoforms, and JNK1a2, JNK1b2, JNK2a2, JNK2b2, and JNK3b2, which represent the p54 isoforms. JNKs coordinate cell responses to stress and influence regulation of cell growth and transformation. The human JNK1 (PRKM8, SAPK1, MAPK8) gene maps to chromosome 10q11.22 and shares 83% amino acid identity with JNK2. JNK1 is necessary for normal activation and differentiation of CD4 helper T (TH) cells into TH1 and TH2 effector cells. Capsaicin activates JNK1 and p38 in ras-transformed human breast epithelial cells. Nitrogen oxides (NOx) upregulate JNK1 in addition to c-Fos, c-Jun, and other signaling kinases, including MEKK1 and p38.

For Reserch Use Only. Not For Use In Diagnostic Procedures