

## PKR Rabbit Monoclonal Antibody

### Catalog #: EAB21454

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC	62	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:10-100
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-200
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-200

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	PKR Rabbit Monoclonal Antibody detects endogenous levels of PKR protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P19525</a>
<b>Entrez-Gene Id</b>	<a href="#">5610</a>

### Product Description

The protein encoded by this gene is a serine/threonine protein kinase that is activated by autophosphorylation after binding to dsRNA. The activated form of the encoded protein can phosphorylate translation initiation factor EIF2S1, which in turn inhibits protein synthesis. This protein is also activated by manganese ions and heparin. The encoded protein plays an important role in the innate immune response against multiple DNA and RNA viruses.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**