

## Cleaved PARP-1 Rabbit Monoclonal Antibody

### Catalog #: EAB21483

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IF/ICC, FC	113	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:10-100
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-200
<b>FC</b> (Flow Cytometry)	1:10-100

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Cleaved PARP-1 Rabbit Monoclonal Antibody detects endogenous levels of Cleaved PARP-1 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P09874</a>
<b>Entrez-Gene Id</b>	<a href="#">142</a>

### Product Description

PARP-1(Poly(ADP-ribose) polymerase-1), also designated PARP, is a nuclear DNA-binding zinc finger protein that influences DNA repair, DNA replication, modulation of chromatin structure, and apoptosis. In response to genotoxic stress, PARP-1 catalyzes the transfer of ADP-ribose units from NAD(+) to a number of acceptor molecules including chromatin. PARP-1 recognizes DNA strand interruptions and can complex with RNA and negatively regulate transcription. Actinomycin D- and etoposide-dependent induction of caspases mediates cleavage of PARP-1 into a p89 fragment that traverses into the cytoplasm. Apoptosis-inducing factor (AIF) translocation from the mitochondria to the nucleus is PARP-1-dependent and is necessary for PARP-1-dependent cell death. PARP-1 deficiencies lead to chromosomal instability due to higher frequencies of chromosome fusions and aneuploidy, suggesting that poly(ADP-ribosyl)ation contributes to the efficient maintenance of genome integrity.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**