

## LC3A Rabbit Monoclonal Antibody

### Catalog #: EAB21533

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC, IF/ICC, FC	14, 16	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:10-100
<b>IHC</b> (Immunohistochemistry)	1:20-200
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:20-200
<b>FC</b> (Flow Cytometry)	1:10-100

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	LC3A Rabbit Monoclonal Antibody detects endogenous levels of LC3A protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">Q9H492</a>
<b>Entrez-Gene Id</b>	<a href="#">84557</a>

### Product Description

Microtubule-associated proteins (MAPs) regulate microtubule stability and play critical roles in neuronal development and in maintaining the balance between neuronal plasticity and rigidity. MAP-light chain 3 beta (MAP-LC3 $\beta$ ) and MAP-light chain 3 alpha (MAP-LC3 $\alpha$ ) are subunits of both MAP1A and MAP1B. MAP-LC3 $\beta$ , a homolog of Apg8p, is essential for autophagy and associated to the autophagosome membranes after processing. Two forms of LC3 $\beta$ , the cytosolic LC3-I and the membrane-bound LC3-II, are produced post-translationally. LC3-I is formed by the removal of the C-terminal 22 amino acids from newly synthesized LC3 $\beta$ , followed by the conversion of a fraction of LC3-I into LC3-II. LC3 enhances fibronectin mRNA translation in ductus arteriosus cells through association with 60S ribosomes and binding to an AU-rich element in the 3' untranslated region of fibronectin mRNA. This facilitates sorting of fibronectin mRNA onto rough endoplasmic reticulum and translation.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**