

Phospho-Tau (Ser396) Rabbit Monoclonal Antibody

Catalog #: EAB21733

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|-------------------|----------|-------------------|
| Rabbit IgG | Monoclonal | WB, IHC-P, IF/ICC | 79 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|------------|
| WB (Western Blotting) | 1:500-2000 |
| IHC-P (Immunohistochemistry-Paraffin) | 1:100-500 |
| IF/ICC (Immunofluorescence/Immunocytochemistry) | 1:50-200 |

Product Information

| | |
|-----------------------|--|
| Conjugate | Unconjugate |
| Specificity | Phospho-Tau (Ser396) Rabbit Monoclonal Antibody detects endogenous levels of tau protein when phosphorylated at Ser396. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | P10636 |
| Entrez-Gene Id | 4137 |

Product Description

Tau, also known as MAPT (microtubule-associated protein tau), MAPTL, MTBT1 or TAU, is a 758 amino acid protein that localizes to the cytoplasm, as well as to the cytoskeleton and the cell membrane, and contains four Tau/MAP repeats. Expressed in neuronal tissue and existing as multiple alternatively spliced isoforms, Tau functions to promote microtubule assembly and stability and is thought to be involved in the maintenance of neuronal polarity. Tau may also link microtubules with neural plasma membrane components and, addition to its role in microtubule stability, is also necessary for cytoskeletal plasticity. Tau is highly subject to a variety of post-translational modifications, including phosphorylation on serine and threonine residues, polyubiquitination (and subsequent proteasomal degradation) and glycation of specific Tau isoforms.

For Reserch Use Only. Not For Use In Diagnostic Procedures