

RIG-I/DDX58 Rabbit Monoclonal Antibody

Catalog #: EAB21900

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB	107	Human, Mouse

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000

Product Information

Conjugate	Unconjugate
Specificity	RIG-I/DDX58 Rabbit Monoclonal Antibody detects endogenous levels of RIG-I/DDX58 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	O95786
Entrez-Gene ID	23586

Product Description

Retinoic acid inducible gene I (RIG-I) is a 925 amino acid, interferon-inducible cellular DExD/H box RNA helicase that activates type I interferon (IFN), an important effector of the innate immune system that is sensitive to these dsRNA viruses. dsRNA is normally present in very low quantities in cells, so when a virus is present, the elevated levels of dsRNA act as a sign telling RIG-I to activate the production of IFN. RIG-I does this by using its helicase domain to bind to viral dsRNA, thus transmitting the activation signal for IFN through I κ B kinase-related kinases and inducing IFN expression. RIG-I is expressed in the cytoplasm of fibroblasts and conventional dendritic cells and can distinguish between many different RNA viruses.

For Reserch Use Only. Not For Use In Diagnostic Procedures