

## NPM Rabbit Monoclonal Antibody

# Catalog #: EAB21980

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P, IF/ICC, FC	37	Human, Mouse, Rat

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:100-500
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-200
FC(Flow Cytometry)	1:10-100

#### **Product Information**

Conjugate	Unconjugate
Specificity	NPM Rabbit Monoclonal Antibody detects endogenous levels of NPM protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>P06748</u>
Entrez-Gene Id	<u>4869</u>

### **Product Description**

The protein encoded by this gene is involved in several cellular processes, including centrosome duplication, protein chaperoning, and cell proliferation. The encoded phosphoprotein shuttles between the nucleolus, nucleus, and cytoplasm, chaperoning ribosomal proteins and core histones from the nucleus to the cytoplasm. This protein is also known to sequester the tumor suppressor ARF in the nucleolus, protecting it from degradation until it is needed. Mutations in this gene are associated with acute myeloid leukemia. Dozens of pseudogenes of this gene have been identified.

For Reserch Use Only. Not For Use In Diagnostic Procedures