

Product Datasheet

Order: order@ebiocell.com

Supprt:

Web:

TEL: (540)808-3925 tech@ebiocell.com

www.ebiocell.com

Tri-Methyl-Histone H3 (Lys27) Rabbit Monoclonal Antibody

Catalog #: EAB22303

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC, IF/ICC, ChIP	15	Human, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IP(Immunoprecipitation)	1:10-100
IHC(Immunohistochemistry)	1:20-200
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:20-200
ChIP(Chromatin Immunoprecipitation)	1:10-100

Product Information

Conjugate Unconjugate

Specificity

Tri-Methyl-Histone H3 (Lys27) Rabbit Monoclonal Antibody detects endogenous levels of

histone H3 only when tri-methylated on Lys27.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P68431

 Entrez-Gene Id
 8350

Product Description

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.